

MTT Cell Proliferation Kit

Colorimetric Plate Readout



OZ Biosciences MTT Cell Proliferation Assay Kit is designed for spectrophotometric quantification of cell growth, viability and proliferation and can be used as a direct indicator of cytotoxicity (such as for screening anticancer drugs) and apoptosis.

Ideal when cells in culture showed abnormalities typical of the Hayflick model -after a long period of normal growth- OZ Biosciences Cellular Senescence Kit provides and easy-to-use and efficient method to determine cellular senesence. During senescence in mammalian cells, an endogenous lysosomal β -galactosidase is over-expressed and is accumulated within the cells. The presence of this β -Gal activity is a marker of aging cell population in vitro.

Cellular Senescence Kit Specific Stem cells Application



Adipose derived Stem Cells (passage 7) and Mesenchymal Stem Cell (passage 6) stained with the Cellular Senescence Kit. Nuclei counterstained with Crystal Violet solution, *not supplied*.

MORE PRODUCTS

D-Luciferin Potassium & Sodium Salt

Highest Purity: 99.7% - Endotoxin Free - Competitive price.

DMNPE-caged Luciferin

A D-Luciferin ester analogue which can cross cell membranes efficiently and can be used to supply a continuous source of active luciferin.

Cas9 Nuclease

Optimized Cas9 Nuclease - with a N-Terminal His Tag, 2 Optimized Nuclear Localization Sequences (NLS) + 1 Genuine Targeting Sequence. Designed to be more efficient at targeting the Cas9 protein to the nucleus.

USA & CANADA

OZ Biosciences USA Inc. 4901 Morena Boulevard, Suite 901 San Diego, CA 92117, USA Ph: +1 858 246 7840 Fax: +1 855 631 0626 E-mail: orderusa@ozbiosciences.com



FRANCE & EUROPE

OZ Biosciences SAS Parc Scientifique de Luminy zone entreprise, case 922 13288 Marseille Cedex 09 FRANCE Ph: +33 486 94 85 16 Fax: +33 486 94 85 15 E-mail: order@ozbiosciences.com



Protein Quantification Assays Enzyme Detection - Reporter Gene Assays Viability / Apoptosis / Stress Assays

Cellular Assay Kits

A COMPREHENSIVE LINE TO EASE AND REFINE YOUR RESEARCH

β-galactosidase & X-Gal Kits

GENE LacZ is one of the most frequently reporter gene used in transfection experiments. LacZ encoded protein, b-galactosidase, is very stable, resistant to proteolytic degradation REPORTER and easily tested.

- Measuring high expression level of β-gal - Choose the ONPG Assay Kit
- Measuring low expression level of β-gal - Choose the CPRG Assay Kit
- Visualization of LacZ transfected cells in vitro and in vivo Choose the X-Gal Staining Kit



The Luciferase Assay Kit is the most sensitive analytical tool for measuring gene expression. Fast and easy, accurate and linear, this kit detects and guantifies firefly luciferase in transfected eukaryotic cells reaching high sensitivity level.

Accurate & Highly Sensitive - Detect as

Simple & Fast - Protocol: 15 min - Pic of signal

Safe & Biodegradable - A natural

product: non toxic, heavy metal free

PICK

THE KIT

THAT BEST

FITS YOUR NEEDS

within 30 min - Signal duration: up to 4 hours

little as 40 ng/mL - Large linear detection

range: Over 3-orders of magnitude

SEAPAssayKit

OZ Biosciences SEAP Assay Kit (Secreted Alkaline Phosphatase) is a colorimetric assay for sensitive quantification of SEAP in culture medium from transfected cells or tissues.

PROTEIN DOSAG

Ē

FluoProdige Protein Assay Kit

Facilitates and Improves Protein and Peptide Quantification

The FluoProdige Protein Quantification

Assay Kit presents a complete fluorometric assay for protein and peptide quantification.

This assay kit uses a stable analogue to epicocconone molecule



that reversibly binds to lysine, arginine and histidine residues in proteins and peptides to yield an intense red-fluorescent product. The fluorescent signal (~518/605) is directly proportional to protein amount among a wide range of protein concentration, rendering this kit highly sensitive.

Bradford PaK

Classic Colorimetric Measurements

BCA PaK

Any auestions?

tech@ozbiosciences.com

The Bradford Pak, ready-to-use kit, is provided with 1X reagent & 2 sets of BSA standards which means that no dilution, filtration or calculation are required. It is based on the binding of Coomassie Brilliant Blue G-250 dye to the proteins and particularly basic and aromatic amino acids residues.

Simply process your assay in a few minutes

The BCA kit is useful for colorimetric detection and quantification of total protein content even in the presence of detergents. It is based on the reduction of Cu2+ to Cu1+ by proteins in alkaline solution. Bicinchoninic acid (BCA) chelates with the reduced copper Cu1+ and forms a water-soluble purple reaction complex that exhibits a strong absorbance at 562 nm.

Robust and detergent-tolerant colorimetric detection

OZBlue Cell Viability Kit (Resazurin)

Check Viability without killing your cells!

STRESS A ready-to-use assay system based on fluorimetric/ colorimetric quantification of metabolic activity in living cells. The Resazurin dye is used as an indicator of cell viability. 1 SENESCENCE - VIABILITY 1: Add 10 µL of OZBlue Solution per well 2: Incubate 30 min to 4h at 37°C 3: Read the signal using a Fluorescent Plate Reader (560 nm Ex / 590 nm En) or a Spectrophotometer (570 nm)

- Non-toxic OZBlue allows additional analysis on proliferating cells (mRNA, cytogenetic, apoptosis, immunophenotyping ...) and continuous monitoring of cultures over time
- **Non-toxic** An improved alternative to [³H]Thymidine incorporation and Tetrazolium reduction Assay (MTT, XTT)
- Simple A single ready-to-use reagent. No need of washing or extraction procedures

Nitric Oxide Detection Kit



The WST-8 Cell Proliferation Assay Kit is a second generation of colormetic assay for the determination of viable cell number and for studying induction of cell proliferation in vitro.

- More sensitive than other formazan-based assays
- Ready to use no solubilization process required
- Colorimetric measurement at 450nm



The Nitirc Oxide Detection Kit is based on the reaction originally described by Griess to detect NO2- in a variety of biological fluids such as cell culture medium, plasma, serum, or urine.

 Highly sensitive: limit of detection ~2μM-125 pmol Detection of NO2- in a linear and stable way Colorimetric measurement at 540-570 nm

